

# Week 12 Discussion

CAS CS 132: Geometric Algorithms

November 20, 2023

During discussion sections, we will go over three problems.

- The first will be a warm-up question, to help you verify your understanding of the material.
- The second will be a solution to a problem on the assignment of the previous week.
- The third will be a problem similar to one on the assignment of the following week.

The remainder of the time will be dedicated to open Q&A.

# 1 Diagonalization by Hand

Diagonalize the following matrix.

$$A = \begin{bmatrix} -3 & -4 \\ 2 & 3 \end{bmatrix}$$

*Solution.* This implies the eigenvalues of this matrix are 1 and  $-1$ .

Next we have to find bases for the eigenspaces of each of these eigenvalues. First, we will find a basis for  $\text{Nul}(A - I)$ . Since

$$A - I = \begin{bmatrix} -4 & -4 \\ 2 & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$$

the solution set of the equation  $(A - I)\mathbf{x} = \mathbf{0}$  can be written as

$$x_2 \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Therefore,  $[-1 \ 1]^T$  forms a basis of the eigenspace of  $A$  for the eigenvalue 1.

For the eigenvalue  $-1$ , we can do the same:

$$A + I = \begin{bmatrix} -2 & -4 \\ 2 & 4 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$$

which implies (by the same process as above) that  $[-2 \ 1]^T$  forms a basis of the eigenspace of  $A$  for  $-1$ . Therefore, we can take

$$P = \begin{bmatrix} -1 & -2 \\ 1 & 1 \end{bmatrix} \quad D = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

Finally, we have to find the inverse of  $P$ . One way to do this is to use the equation

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

In this case,

$$P^{-1} = \frac{1}{(-1)(1) - (-2)(1)} \begin{bmatrix} 1 & 2 \\ -1 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -1 & -1 \end{bmatrix}$$

## 2 Determinant

Consider the following reduction sequence

$$\begin{aligned}R_1 &\leftarrow R_1 + R_2 \\ \text{swap}(R_2, R_3) \\ R_3 &\leftarrow R_3 + 5R_4 \\ R_2 &\leftarrow -3R_2 \\ R_5 &\leftarrow R_5 - 10R_3 \\ R_5 &\leftarrow R_5/11 \\ \text{swap}(R_5, R_3) \\ \text{swap}(R_1, R_2) \\ R_4 &\leftarrow R_4 + R_1 \\ R_2 &\leftarrow 5R_2 \\ R_1 &\leftarrow -R_1\end{aligned}$$

Suppose that  $A \in \mathbb{R}^{5 \times 5}$  reduces to  $U$  by this reduction sequence, where  $U$  is in *reduced* echelon form. If  $\text{rank } A = 5$ , what is  $\det A$ ?

*Solution.*

### 3 $2 \times 2$ Triangular Matrices and Diagonalization

Consider an arbitrary  $2 \times 2$  matrix

$$A = \begin{bmatrix} a & b \\ 0 & d \end{bmatrix}$$

- A. Find an expression for the characteristic polynomial of  $A$ .
- B. If  $a \neq d$ , then  $A$  has 2 distinct eigenvalues, and so there must be an eigenbasis of  $\mathbb{R}^2$  for  $A$ . If  $a = d$ , for what values of  $b$  is there an eigenbasis of  $\mathbb{R}^2$  for  $A$ ? Justify your answer.

*Solution.*