

# Matrices of Linear Transformations

**Geometric Algorithms**

**Lecture 9**

# Practice Problem

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = 9 \qquad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = 2$$

*Suppose that  $T$  is a linear transformation with the above input-output behavior.*

*What is the domain of  $T$ ? What is the codomain of  $T$ ?*

***What is the value of  $T\left(\begin{bmatrix} 2 \\ -3 \end{bmatrix}\right)$ ?***

**Answer**

domain:  $\mathbb{R}^2$   
codomain:  $\mathbb{R}$

$$T\left(\begin{bmatrix} 2 \\ -3 \end{bmatrix}\right)$$

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = 9 \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = 2$$

$$\begin{bmatrix} 2 \\ -3 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + (-3) \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 2 \\ -3 \end{bmatrix}\right) = T\left(2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + -3 \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$$

$$= 2 T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + -3 T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = 18 - 6 = \boxed{12}$$

# Objectives

1. Look at more examples of linear transformations
2. Show that matrix transformations and linear transformations are really the same thing
3. See more the geometry of linear transformations
4. Relate the properties of matrix equations to properties of linear transformations

# Keywords

matrix of a linear transformation

standard basis vectors (standard coordinate vectors)

2D linear transformations

the unit square

one-to-one

onto

**Recap**

# Recall: Matrices as Transformations

Matrices allow us to *transform* vectors.

The transformed vector lies in the span of its columns.

$$\mathbf{x} \mapsto A\mathbf{x}$$

map a vector  $\mathbf{x}$  to the vector  $A\mathbf{x}$

# Recall: Transformation of a Matrix

The **transformation of a**  $(m \times n)$  **matrix**  $A$  is the function  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  such that

$$T(\mathbf{v}) = A\mathbf{v}$$

given  $\mathbf{v}$ , return  $A$  multiplied by  $\mathbf{v}$

e.g.  $T(\mathbf{v}) = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \mathbf{v}$

$$\begin{aligned} T \left( \begin{bmatrix} 2 \\ 3 \end{bmatrix} \right) &= \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \end{bmatrix} \\ &= 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 3 \end{bmatrix} \end{aligned}$$



# Recall: Motivating Questions

What kind of functions can we define in this way?

How do we interpret what the transformation does to a set of vectors?

How does this relate back to matrix equations?

# Recall: Linear Transformations

**Definition.** A transformation  $T: \mathbb{R}^m \rightarrow \mathbb{R}^n$  is *linear* if it satisfies the following two properties.

1.  $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$  (additivity)

2.  $T(c\mathbf{v}) = cT(\mathbf{v})$  (homogeneity)

# Recall: Linear Transformations

**Definition.** A transformation  $T: \mathbb{R}^m \rightarrow \mathbb{R}^n$  is *linear* if it satisfies the following two properties.

1.  $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$       (additivity)

2.  $T(c\mathbf{v}) = cT(\mathbf{v})$       (homogeneity)

Matrix transformations are linear transformations.

# Properties of Linear Transformations

# The Zero Vector

$$T(\mathbf{0}) = ???$$

# The Zero Vector

$$T(\mathbf{0}) = \mathbf{0}$$

# The Zero Vector

$$T(\mathbf{0}) = \mathbf{0}$$

The zero vector is *fixed* by linear transformations.

# The Zero Vector

$$T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$$

$$T(\mathbf{0}) = \mathbf{0}$$

$$T\left(\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Note: These may be different dimensions!

The zero vector is *fixed* by linear transformations.

$$\begin{aligned} T(\vec{0}) &= T(0\vec{v}) \\ &= 0T(\vec{v}) \\ &= \vec{0} \end{aligned}$$



# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

We can combine our linearity conditions:

# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

We can combine our linearity conditions:

$$T(a\mathbf{v} + b\mathbf{u})$$

# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

We can combine our linearity conditions:

$$\begin{aligned} T(a\mathbf{v} + b\mathbf{u}) \\ = T(a\mathbf{v}) + T(b\mathbf{u}) \quad (\text{additivity}) \end{aligned}$$

# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

We can combine our linearity conditions:

$$T(a\mathbf{v} + b\mathbf{u})$$

$$= T(a\mathbf{v}) + T(b\mathbf{u}) \quad (\text{additivity})$$

$$= aT(\mathbf{v}) + bT(\mathbf{u}) \quad (\text{homogeneity for each term})$$

# A Single Condition

**Theorem.** A transformation  $T: \mathbb{R}^m \rightarrow \mathbb{R}^n$  is linear if and only if for any vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathbb{R}^m$  and any real numbers  $a$  and  $b$ ,

$$T(a\mathbf{u} + b\mathbf{v}) = aT(\mathbf{u}) + bT(\mathbf{v})$$

# A Single Condition

**Theorem.** A transformation  $T: \mathbb{R}^m \rightarrow \mathbb{R}^n$  is linear if and only if for any vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathbb{R}^m$  and any real numbers  $a$  and  $b$ ,

$$T(a\mathbf{u} + b\mathbf{v}) = aT(\mathbf{u}) + bT(\mathbf{v})$$

It's often easiest to show this single condition.

# Linear Combinations

$$T(a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n) = a_1T(\mathbf{v}_1) + a_2T(\mathbf{v}_2) + \dots + a_nT(\mathbf{v}_n)$$

We can generalize this condition to any linear combination.



# Linear Combinations

$$T\left(\sum_{i=1}^n a_i \mathbf{v}_i\right) = \sum_{i=1}^n a_i T(\mathbf{v}_i)$$

We can generalize this condition to any linear combination.

# Linear Combinations

$$T\left(\sum_{i=1}^n a_i \mathbf{v}_i\right) = \sum_{i=1}^n a_i T(\mathbf{v}_i)$$

We can generalize this condition to any linear combination.

This is the most useful form.

# Example: Identity

$$T(\mathbf{v}) = \mathbf{v}$$


$$T(\vec{v} + \vec{w}) = \vec{v} + \vec{w} \quad \text{(additivity)} \quad \checkmark$$

$$= T(\vec{v}) + T(\vec{w})$$

$$T(c\vec{v}) = c\vec{v} = cT(\vec{v}) \quad \text{(homogeneity)} \quad \checkmark$$



# Example: Zero

$$T(\mathbf{v}) = \mathbf{0}$$

$$\begin{aligned} T(a\vec{v}_1 + b\vec{v}_2) &= \mathbf{0} \\ &= \mathbf{0} + \mathbf{0} \\ &= a\mathbf{0} + b\mathbf{0} \\ &= aT(\vec{v}_1) + bT(\vec{v}_2) \end{aligned}$$


# Example: Rotation

We'll see this on Thursday, but we can reason about it geometrically for now.



# Non-Example: Squares

$$T(x) = x^2$$

Note that  $T: \mathbb{R}^1 \rightarrow \mathbb{R}^1$

$$T(5(1)) = 25$$

$$5 T(1) = 5$$


does not satisfy homogeneity

# Non-Example: Translation


$$T(2 \begin{bmatrix} 1 \\ 1 \end{bmatrix}) = \begin{bmatrix} 3 \\ 3 \end{bmatrix}$$

$$2 T(\begin{bmatrix} 1 \\ 1 \end{bmatrix}) = \begin{bmatrix} 4 \\ 4 \end{bmatrix}$$

$$2 (\begin{bmatrix} 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 1 \end{bmatrix}) = 2 \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \end{bmatrix}$$



$$\mathbf{x} + \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



# Example (Understanding Check)

$$T(\mathbf{v}) = 5\mathbf{v}$$

$$\begin{aligned} T(a\vec{v} + b\vec{u}) &= 5(a\vec{v} + b\vec{u}) \\ &= 5a\vec{v} + 5b\vec{u} \\ &= a5\vec{v} + b5\vec{u} \\ &= aT(\vec{v}) + bT(\vec{u}) \end{aligned}$$



# Example (Understanding Check)

$$T(x) = e^x$$

$$T(0) = 1$$

# Properties of Linear Transformations

# The Zero Vector

$$T(\mathbf{0}) = ???$$

# The Zero Vector


$$T(\mathbf{0}) = \mathbf{0}$$

# The Zero Vector

$$T(\mathbf{0}) = \mathbf{0}$$

The zero vector is *fixed* by linear transformations.  
It can't move anywhere.

# The Zero Vector

$$T(\mathbf{0}) = \mathbf{0}$$


Note: These may be different dimensions!

The zero vector is *fixed* by linear transformations.  
It can't move anywhere.

# Verification

any matrix transformation:

rotation about the origin:

translation (*non-example*):

# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$



# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

We can combine our linearity conditions:

# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

We can combine our linearity conditions:

$$T(a\mathbf{v} + b\mathbf{u})$$

# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

We can combine our linearity conditions:

$$T(a\mathbf{v} + b\mathbf{u})$$

$$= T(a\mathbf{v}) + T(b\mathbf{u}) \quad (\text{by additivity})$$

# A Single Condition

$$T(a\mathbf{v} + b\mathbf{u}) = aT(\mathbf{v}) + bT(\mathbf{u})$$

We can combine our linearity conditions:

$$T(a\mathbf{v} + b\mathbf{u})$$

$$= T(a\mathbf{v}) + T(b\mathbf{u}) \quad (\text{by additivity})$$

$$= aT(\mathbf{v}) + bT(\mathbf{u}) \quad (\text{by homogeneity for each term})$$

# A Single Condition

**Theorem.** A transformation  $T: \mathbb{R}^m \rightarrow \mathbb{R}^n$  is linear if and only if for any vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathbb{R}^m$  and any real numbers  $a$  and  $b$ ,

$$T(a\mathbf{u} + b\mathbf{v}) = aT(\mathbf{u}) + bT(\mathbf{v})$$

## Example (Again)

$$T(\mathbf{v}) = 5\mathbf{v}$$

# Linear Combinations

$$T(a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n) = a_1T(\mathbf{v}_1) + a_2T(\mathbf{v}_2) + \dots + a_nT(\mathbf{v}_n)$$

We can generalize this condition to any linear combination.

# Linear Combinations

$$T\left(\sum_{i=1}^n a_i \mathbf{v}_i\right) = \sum_{i=1}^n a_i T(\mathbf{v}_i)$$

We can generalize this condition to any linear combination.



# Linear Combinations

$$T\left(\sum_{i=1}^n a_i \mathbf{v}_i\right) = \sum_{i=1}^n a_i T(\mathbf{v}_i)$$

We can generalize this condition to any linear combination.

This is the most useful form.

# **Our Next Motivating Question**

# Our Next Motivating Question

We know that matrix transformations are linear transformations.

# Our Next Motivating Question

We know that matrix transformations are linear transformations.

*Are there any other kinds of linear transformations?*

# Our Next Motivating Question

We know that matrix transformations are linear transformations.

*Are there any other kinds of linear transformations?*

No

# Matrix of a Linear Transformation

**Theorem.** A transformation  $T$  is linear if and only if there is a matrix whose corresponding transformation is  $T$  (which "implements"  $T$ ).

# Matrix of a Linear Transformation

**Theorem.** A transformation  $T$  is linear if and only if there is a matrix whose corresponding transformation is  $T$  (which "implements"  $T$ ).

Linear transformations are **exactly**  
matrix transformations.

# A Fundamental Concern

*Given a linear transformation  $T$ , how do we find the matrix  $A$  such that*

$$T(\mathbf{v}) = A\mathbf{v}?$$



# A Thought Experiment

Suppose I tell you  $T$  is a linear transformation and

$$T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 3 \\ 4 \end{bmatrix} \quad T\left(\begin{bmatrix} 3 \\ 4 \end{bmatrix}\right) = \begin{bmatrix} 5 \\ 6 \end{bmatrix}$$

Do we know what  $T\left(\begin{bmatrix} 4 \\ 6 \end{bmatrix}\right)$  is?

**Answer: Yes**

$$T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 3 \\ 4 \end{bmatrix}\right) = \begin{bmatrix} 5 \\ 6 \end{bmatrix}$$

Because of additivity:

$$\begin{aligned} T\left(\begin{bmatrix} 4 \\ 6 \end{bmatrix}\right) &= T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix} + \begin{bmatrix} 3 \\ 4 \end{bmatrix}\right) \\ &= T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) + T\left(\begin{bmatrix} 3 \\ 4 \end{bmatrix}\right) = \\ &\quad \begin{bmatrix} 3 \\ 4 \end{bmatrix} + \begin{bmatrix} 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 8 \\ 10 \end{bmatrix} \end{aligned}$$

# A Thought Experiment

$$T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 3 \\ 4 \end{bmatrix} \quad T\left(\begin{bmatrix} 3 \\ 4 \end{bmatrix}\right) = \begin{bmatrix} 5 \\ 6 \end{bmatrix}$$

What about:

$$T\left(\begin{bmatrix} 2 \\ 3 \end{bmatrix}\right) = T\left(\frac{1}{2}\begin{bmatrix} 4 \\ 6 \end{bmatrix}\right) = \frac{1}{2}T\left(\begin{bmatrix} 4 \\ 6 \end{bmatrix}\right)$$

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 3 \\ 4 \end{bmatrix} - a\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

# The Takeaway

Linearity is a **very** strong restriction.

If we know the values of  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  on **any** set of vectors which spans all of  $\mathbb{R}^n$ , then we know  $T$ .

why? :

# **Another Thought Experiment (Game)**

# Another Thought Experiment (Game)

Suppose I am holding a matrix  $A$ .

# Another Thought Experiment (Game)

Suppose I am holding a matrix  $A$ .

Your objective is to figure out what  $A$  is.

# Another Thought Experiment (Game)

Suppose I am holding a matrix  $A$ .

Your objective is to figure out what  $A$  is.

But you're only allowed to ask the question:



# Another Thought Experiment (Game)

Suppose I am holding a matrix  $A$ .

Your objective is to figure out what  $A$  is.

But you're only allowed to ask the question:

what is  $Av$ ?

# Another Thought Experiment (Game)

Suppose I am holding a matrix  $A$ .

Your objective is to figure out what  $A$  is.

But you're only allowed to ask the question:

what is  $Av$ ?

(you pick the  $v$ 's, and I have to tell the truth)

# Another Thought Experiment (Game)

Suppose I am holding a matrix  $A$ .

Your objective is to figure out what  $A$  is.

But you're only allowed to ask the question:

what is  $Av$ ?

(you pick the  $v$ 's, and I have to tell the truth)

This is basically linear algebraic battleship.

# Recall: Calculating $A\mathbf{v}$

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ \vdots \\ ? \end{bmatrix}$$

$$b_1 = a_{11}v_1 + a_{12}v_2 + \cdots + a_{1n}v_n = \sum_{i=1}^n a_{1i}v_i$$

# Recall: Matrix-Vector Multiplication

**Definition.** Given a  $(m \times n)$  matrix  $A$  with columns  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$ , and a vector  $\mathbf{v}$  in  $\mathbb{R}^n$ , we define

$$A\mathbf{v} = [\mathbf{a}_1 \quad \mathbf{a}_2 \quad \dots \quad \mathbf{a}_n] \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = v_1\mathbf{a}_1 + v_2\mathbf{a}_2 + \dots + v_n\mathbf{a}_n$$

# Recall: Matrix-Vector Multiplication

**Definition.** Given a  $(m \times n)$  matrix  $A$  with columns  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$ , and a vector  $\mathbf{v}$  in  $\mathbb{R}^n$ , we define

$$A\mathbf{v} = [\mathbf{a}_1 \quad \mathbf{a}_2 \quad \dots \quad \mathbf{a}_n] \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = v_1\mathbf{a}_1 + v_2\mathbf{a}_2 + \dots + v_n\mathbf{a}_n$$

# Recall: Matrix-Vector Multiplication

**Definition.** Given a  $(m \times n)$  matrix  $A$  with columns  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$ , and a vector  $\mathbf{v}$  in  $\mathbb{R}^n$ , we define

$$A\mathbf{v} = [\mathbf{a}_1 \quad \mathbf{a}_2 \quad \dots \quad \mathbf{a}_n] \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = v_1\mathbf{a}_1 + v_2\mathbf{a}_2 + \dots + v_n\mathbf{a}_n$$

$A\mathbf{v}$  is a linear combination of the columns of  $A$  with weights given by  $\mathbf{v}$

**Isolating**  $a_{11}$

$$b_1 = a_{11}v_1 + a_{12}v_2 + \dots + a_{1n}v_n = \sum_{i=1}^n a_{1i}v_i$$



**Isolating**  $a_{11}$

$$b_1 = a_{11}v_1 + a_{12}v_2 + \dots + a_{1n}v_n = \sum_{i=1}^n a_{1i}v_i$$

We actually get the whole column  $\mathbf{a}_1$

So its like battleship, but you get to choose one column at a time.

# The Takeaway

We can learn the first column of the matrix implementing

$$T \text{ by looking at } T \begin{pmatrix} \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} \end{pmatrix}$$

# Matrix of a Linear Transformation

# Standard Basis

**Definition.** The *n-dimensional standard basis vectors* (or standard coordinate vectors) are the vectors  $\mathbf{e}_1, \dots, \mathbf{e}_n$  where

$$\mathbf{e}_i = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix} \begin{matrix} 1 \\ 2 \\ \vdots \\ i-1 \\ i \\ i+1 \\ \vdots \\ n-1 \\ n \end{matrix}$$

# Standard Basis

**Definition (Alternative).** The  $n$ -dimensional standard basis vectors  $\mathbf{e}_1, \dots, \mathbf{e}_n$  are the columns of the  $n \times n$  identity matrix.

$$I = [\mathbf{e}_1 \quad \mathbf{e}_2 \quad \dots \quad \mathbf{e}_n]$$

# Standard Basis and the Matrix Equation

The key points:  $[\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n] \mathbf{e}_i = \mathbf{a}_i$

The standard basis vectors gives us a way to "look into" a matrix.

# Standard Basis and Vector Coordinates

$$\begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = v_1 \mathbf{e}_1 + v_2 \mathbf{e}_2 + \dots + v_n \mathbf{e}_n$$

Column vectors can be viewed as describing how to write a vector as a linear combination of the standard basis.

Example:

# Standard Basis and Linear Transformations

**Theorem.** For any linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ , the matrix

$$A = [T(\mathbf{e}_1) \quad T(\mathbf{e}_2) \quad \dots \quad T(\mathbf{e}_n)]$$

is the unique matrix such that  $T(\mathbf{v}) = A\mathbf{v}$  for all  $\mathbf{v}$  in  $\mathbb{R}^n$ .



# More Formally

$$T(\mathbf{v}) =$$

$$= \begin{bmatrix} T(\mathbf{e}_1) & T(\mathbf{e}_2) & \dots & T(\mathbf{e}_n) \end{bmatrix} \mathbf{v}$$

# How To: Matrices of Linear Transformations

**Question.** Find the matrix which implements the transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ .

**Solution.** Determine the images of standard basis under  $T$ . Then write down

$$[T(\mathbf{e}_1) \quad T(\mathbf{e}_2) \quad \dots \quad T(\mathbf{e}_n)]$$


# Question

*Write down the matrix which implements the linear transformation  $T$  which **rotates** vectors by 90 degrees clockwise.*


**Answer**

# General Rotation

Additivity



Homogeneity



Rotation a Linear Transformation

# Geometry of Matrix Transformations

# Motivating Questions

What kind of functions can we define in this way?

How do we interpret what the transformation does to a set of vectors?



How does this relate back to matrix equations?

# Motto

Matrix transformations change the "shape" of a set of set of vectors (points).




# Example: Dilation





# Example: Dilation

$$\begin{bmatrix} r & 0 \\ 0 & r \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} rx_1 \\ rx_2 \end{bmatrix}$$




if  $r > 1$ , then the transformation pushes points away from the origin.

# Example: Contraction



# Example: Contraction

$$\begin{bmatrix} r & 0 \\ 0 & r \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} rx_1 \\ rx_2 \end{bmatrix}$$




if  $0 \leq r \leq 1$ , then the transformation pulls points towards the origin.

# Example: Shearing




# Example: Shearing

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 + x_2 \\ x_2 \end{bmatrix}$$




$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \mathbf{x}$$




Imagine shearing like with rocks or metal.

# Question




Draw how this matrix transforms points. What kind of transformation does it represent?

# Answer: Reflection




$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{x}$$





# General Rotation

How does  
rotation  
affect the  
standard  
basis?



# Rotation Matrix

# Rotation Matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

# Rotation Matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

**Note:** This is rotation about the origin.

# Rotation Matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

**Note:** This is rotation about the origin.


**The Takeaway:** We can figure out the matrices which implement complex linear transformations by understanding what they do to the standard basis.

# Question (Conceptual)

*Is rotation about a point other than the origin a linear transformation?*


**Answer: No**

The origin is not  
fixed by this  
transformation.



# The Unit Square

The *unit square* is the set of points in  $\mathbb{R}^2$  enclosed by the points  $(0,0)$ ,  $(0,1)$ ,  $(1,0)$ ,  $(1,1)$ .





# How To: The Unit Square and Matrices

# How To: The Unit Square and Matrices

**Question.** Find the matrix which implements the linear transformation which is represented geometrically in the following picture.


# How To: The Unit Square and Matrices

**Question.** Find the matrix which implements the linear transformation which is represented geometrically in the following picture.


**Solution.** Find where the standard basis vectors go.

# Question

*Write down the matrix for the following shearing operation using this method.*



# Answer




You need to **know** these matrices, but you don't need to memorize them.

**Remember:** What does this matrix do to the unit square? Then build the matrix from there.


# Reflection through the $x_2$ -axis

Reflection through the  $x_2$  axis





# Projections

Projection onto the  $x_1$  axis





# A 3D Example: Rotation about the $x_3$ -Axis ( $z$ -Axis)



# List of Important 2D Linear Transformations

- » dilation, contraction
- » reflections
- » projections
- » horizontal/vertical contractions
- » horizontal/vertical shearing

Look through the notes for a comprehensive collection of pictures or...

demo

# One-to-One and Onto

# Recall: Motivating Questions

What kind of functions can we define in this way?

How do we interpret what the transformation does to a set of vectors?

How does this relate back to matrix equations?

# Recall: A New Interpretation of the Matrix Equation

$A\mathbf{x} = \mathbf{b}$ ?  $\equiv$  is there a vector which  $A$   
transforms into  $\mathbf{b}$ ?

Solve  $A\mathbf{x} = \mathbf{b}$   $\equiv$  find a vector which  $A$   
transforms into  $\mathbf{b}$

# Recall: A New Interpretation of the Matrix Equation

$A\mathbf{x} = \mathbf{b}$ ?  $\equiv$  is there a vector which  $A$   
transforms into  $\mathbf{b}$ ?

Solve  $A\mathbf{x} = \mathbf{b}$   $\equiv$  find a vector which  $A$   
transforms into  $\mathbf{b}$

What about other questions?

# Other Questions Like...

Does  $A\mathbf{x} = \mathbf{b}$  have a solution for any choice of  $\mathbf{b}$ ?

Does  $A\mathbf{x} = \mathbf{0}$  have a unique solution?



# Other Questions Like...

Does  $A\mathbf{x} = \mathbf{b}$  have **at least one solution** for any choice of  $\mathbf{b}$ ?

Does  $A\mathbf{x} = \mathbf{b}$  have **at most one solution** for any choice of  $\mathbf{b}$ ?

# Wait

$A\mathbf{x} = \mathbf{0}$  has a  
unique solution

$\equiv$

$A\mathbf{x} = \mathbf{b}$  has at most one  
solution


why? :

# Onto and One-to-One


**Definition.** A transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is **onto** if any vector  $\mathbf{b}$  in  $\mathbb{R}^m$  is the **image of at least one vector**  $\mathbf{v}$  in  $\mathbb{R}^n$  (where  $T(\mathbf{v}) = \mathbf{b}$ ).

**Definition.** A transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is **one-to-one** if any vector  $\mathbf{b}$  in  $\mathbb{R}^m$  is the **image of at most one vector**  $\mathbf{v}$  in  $\mathbb{R}^n$  (where  $T(\mathbf{v}) = \mathbf{b}$ ).

# Onto (Pictorially)




$T$  is *not* onto  $\mathbb{R}^m$




$T$  is onto  $\mathbb{R}^m$

# Onto (Pictorially)




$T$  is *not* onto  $\mathbb{R}^m$




$T$  is onto  $\mathbb{R}^m$

$T$  is onto if its range = its codomain

# One-to-One (Pictorially)



$T$  is *not* one-to-one



$T$  is one-to-one

# Taking Stock: Onto

**Theorem.** The following are logically equivalent for the linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  implemented by the matrix  $A$ .

- »  $T$  is onto
- »  $A\mathbf{x} = \mathbf{b}$  has a solution for any choice of  $\mathbf{b}$
- »  $\text{range}(T) = \text{codomain}(T)$
- » the columns of  $A$  span  $\mathbb{R}^m$
- »  $A$  has a pivot position in every row

# Taking Stock: One-to-One

**Theorem.** The following are logically equivalent for the linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  implemented by the matrix  $A$ .

- »  $T$  is one-to-one
- »  $A\mathbf{x} = \mathbf{b}$  has at most one solution for any  $\mathbf{b}$
- »  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution
- » The columns of  $A$  are linearly independent
- »  $A$  has a pivot position in every column



# How To: One-to-One and Onto

**Question.** Show that the linear transformation  $T$  is one-to-one/onto.

**Solution.** (one approach) Find the matrix which implements  $T$  and see if it has a pivot in every column/row.

Warning: this is not the only way. Always try to think if you can solve it using *any* of the perspectives

# Example: both 1-1 and onto

Rotation about the origin:

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

why? :

# Example: 1-1, not onto

Lifting:

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \mapsto \begin{bmatrix} x_1 \\ x_2 \\ x_1 + x_2 \end{bmatrix}$$


why? :

# Example: not 1-1, not onto

Projection onto the  $x_1$  axis:

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

why? :




# Example: onto, not 1-1

Projection from  $\mathbb{R}^3$  to  $\mathbb{R}^2$ .

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \mapsto \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

why? :



# Summary

Matrix transformations and linear transformations are the same thing.

We can find these matrices by looking at how the transformation behaves on the standard basis.

We can reason about matrix equations by directly reasoning about the linear transformations.