# LU Factorization

Geometric Algorithms
Lecture 12

#### Practice Problem

 1
 2

 3
 7

Determine the inverse of the above matrix in every way that we've discussed

## nswer

$$\begin{bmatrix} a & b \end{bmatrix}^{-1} = \frac{1}{ad - bd} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & 1 \\ 3 & 7 \end{bmatrix}^{-1} = \frac{1}{7 - 6} \begin{bmatrix} 7 & -2 \\ -3 & 1 \end{bmatrix}^{-2} \begin{bmatrix} 7 & -2 \\ -3 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 7 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -3 \end{bmatrix}$$

\[ \left( \text{1 0 7} \) \[ \text{0 1 -3} \]

### Answer

$$\begin{bmatrix}
1 & 2 & 1 & 0 \\
3 & 7 & 0 & 0
\end{bmatrix} \sim \begin{bmatrix}
1 & 2 & 1 & 0 \\
0 & 1 & -3 & 1
\end{bmatrix} \sim \begin{bmatrix}
1 & 0 & 7 & -2 \\
0 & 1 & -3 & 1
\end{bmatrix}$$

$$A \hat{x} = \begin{bmatrix} 3 \\ 10 \end{bmatrix} \begin{bmatrix} \frac{1}{3} \cdot \frac{1}{2} \cdot \frac{3}{10} \\ \frac{3}{7} \cdot \frac{1}{10} \end{bmatrix} \sim \begin{bmatrix}
1 & 0 & 1 \\
0 & 1 & 1
\end{bmatrix} \sim \begin{bmatrix}
1 & 0 & 1 \\
0 & 1 & 1
\end{bmatrix}$$

$$x = \begin{pmatrix} 7 & -7 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 10 \end{pmatrix} = \begin{pmatrix} 21 & -20 \\ -9 & +10 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ \end{bmatrix}$$

## Objectives

- » Motivate matrix factorization in general, and the LU factorization in specific
- » Recall elementary row operations and connect
  them to matrices
- » Look at the LU factorization, how to find it, and how to use it

## LU Factorization

## Matrix Factorization

#### Matrix Factorization

A **factorization** of a matrix A is an equation which expresses A as a product of one or more matrices, e.g.,

$$A = BC$$

#### Matrix Factorization

A **factorization** of a matrix A is an equation which expresses A as a product of one or more matrices, e.g.,

$$A = BC$$

So far, we've been given two factors and asked to find their product

Factorization is the harder direction

Writing A as the product of multiple matrices can

Writing A as the product of multiple matrices can

 $\gg$  make computing with A faster

Writing A as the product of multiple matrices can

 $\gg$  make computing with A faster

 $\gg$  make working with A easier

Writing A as the product of multiple matrices can

- >> make computing with A faster
- $\gg$  make working with A easier
- $\gg$  expose important information about A

Writing A as the product of multiple matrices can

- » make computing with A faster LU Decomposition
- $\gg$  make working with A easier
- $\gg$  expose important information about A

Question. For an matrix A, solve the equations

$$A\mathbf{x} = \mathbf{b}_1$$
 ,  $A\mathbf{x} = \mathbf{b}_2$  ...  $A\mathbf{x} = \mathbf{b}_{k-1}$  ,  $A\mathbf{x} = \mathbf{b}_k$ 

In other words: we want to solve <u>several</u> matrix equations over the same matrix

**Question.** For a matrix A, solve (for X) in the equation

$$AX = B$$

where X and B are matrices of appropriate dimension

This is (essentially) the same question

Question. Solve AX = B

If A is invertible, then we have a solution:

Find  $A^{-1}$  and then  $X = A^{-1}B$ 

Question. Solve AX = B

If A is invertible, then we have a solution:

Find  $A^{-1}$  and then  $X = A^{-1}B$ 

What if  $A^{-1}$  is not invertible? Even if it is, can we do it faster?

## LU Factorization at a High Level

Given a  $m \times n$  matrix A, we are going to factorize A as

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ * & 1 & 0 & 0 \\ * & * & 1 & 0 \\ * & * & * & 1 \end{bmatrix} \begin{bmatrix} \blacksquare & * & * & * & * \\ 0 & \blacksquare & * & * & * \\ 0 & 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$L \qquad U$$

## LU Factorization at a High Level

Given a  $m \times n$  matrix A, we are going to factorize A as

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ * & 1 & 0 & 0 \\ * & * & 1 & 0 \\ * & * & * & 1 \end{bmatrix} \begin{bmatrix} \blacksquare & * & * & * & * \\ 0 & \blacksquare & * & * & * \\ 0 & 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$L \qquad U$$

Note. This applies to non-square matrices

#### What are "L" and "U"?

L stands for "lower" as in *lower triangular*U stands for "upper" as in *upper triangular* 

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ * & 1 & 0 & 0 \\ * & * & 1 & 0 \\ * & * & * & 1 \end{bmatrix} \begin{bmatrix} \bullet & * & * & * \\ 0 & \bullet & * & * \\ 0 & 0 & \bullet & * \\ 0 & 0 & \bullet & \bullet \end{bmatrix}$$

$$L \qquad U$$

$$A = LU$$
 echelon form of  $A$ 

We know how to build U, that's just the forward phase of Gaussian elimination

$$A = LU$$
 echelon form of  $A$ 

We know how to build U, that's just the forward phase of Gaussian elimination

How do we build L?

$$EA = U$$

$$A = LU \text{ echelon form of } A$$

We know how to build U, that's just the forward phase of Gaussian elimination

How do we build L?

The idea. L "implements" the row operations of the forward phase (in rows)

# Elementary Matrices

## Recall: Elementary Row Operations

scaling multiply a row by a number

interchange switch two rows

replacement add a scaled equation to another

## The First Key Observation

## The First Key Observation

Elementary row operations are linear transformations (viewed as transformation on columns)

## The First Key Observation

Elementary row operations are linear transformations (viewed as transformation on columns)

Example: Scale row 2 by 5

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \quad R_2 \leftarrow 5R_2 \quad \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 5a_{21} & 5a_{22} & 5a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

## Example: Scaling

Restricted to one column, we see this is the above linear transformation

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \mapsto \begin{bmatrix} v_1 \\ 5v_2 \\ v_3 \end{bmatrix}$$

## Example: Scaling

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \mapsto \begin{bmatrix} v_1 \\ 5v_2 \\ v_3 \end{bmatrix}$$

Let's build the matrix which implements it:

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 3 \\ 10 & 15 & 30 \\ 7 & 8 & 9 \end{bmatrix}$$

## Another Example: Scaling + Replacement

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \longrightarrow \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ (a_{31} - 2a_{11}) & (a_{32} - 2a_{12}) & (a_{33} - 2a_{13}) \end{bmatrix}$$

$$R_3 \leftarrow (R_3 - 2R_1)$$

## Another Example: Scaling + Replacement

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \longrightarrow \begin{bmatrix} x_1 \\ x_2 \\ x_3 - 2x_1 \end{bmatrix}$$

$$R_3 \leftarrow (R_3 - 2R_1)$$

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \mapsto \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

# Elementary row operations are linear, so they are implemented by matrices

## General Elementary Scaling Matrix

```
\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & k & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}
```

# General Elementary Scaling Matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & k & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

If we want to perform  $R_3 \leftarrow kR_3$  then we need the identity matrix but with the entry  $A_{33} = k$ .

# **General Elementary Scaling Matrix**

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & k & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

If we want to perform  $R_3 \leftarrow kR_3$  then we need the identity matrix but with the entry  $A_{33} = k$ .

If we want to perform  $R_i \leftarrow kR_i$  then we need the identity matrix but with then entry  $A_{ii} = k$ .

#### General Replacement Matrix

```
\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ k & 0 & 0 & 1 \end{bmatrix}
```

#### General Replacement Matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ k & 0 & 0 & 1 \end{bmatrix}$$

If we want to perform  $R_4 \leftarrow R_4 + kR_1$ , then we need the identity matrix but with the entry  $A_{41} = k$ .

#### General Replacement Matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ k & 0 & 0 & 1 \end{bmatrix}$$

If we want to perform  $R_4 \leftarrow R_4 + kR_1$ , then we need the identity matrix but with the entry  $A_{41} = k$ .

If we want to perform  $R_i \leftarrow R_i + kR_j$ , then we need the identity matrix but with the entry  $A_{ij} = k$ .

#### General Swap Matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

If we want to swap  $R_2$  and  $R_3$ , then we need the identity matrix, but with  $R_2$  and  $R_3$  swapped.

# Elementary Matrices

Definition. An elementary matrix is a matrix obtained by applying a single row operation to the identity matrix I

Example. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \xrightarrow{F_1 \leftarrow F_2 + 3F_1} \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

## How To: Finding Elementary Matrices

**Question.** Find the matrix implementing the elementary row operation op

**Solution.** Apply op to the identity matrix of the appropriate size

Taking stock:

#### Taking stock:

» Elementary matrices implement elementary row operations

#### Taking stock:

- » Elementary matrices implement elementary row operations
- » Remember that Matrix multiplication is transformation composition (i.e., do one then the other)

#### Taking stock:

- » Elementary matrices implement elementary row operations
- » Remember that Matrix multiplication is transformation composition (i.e., do one then the other)

So we can implement <u>any</u> sequence of row operations as a product of elementary matrices

#### How to: Matrices implementing Row Operations

**Question.** Find the matrix implementing a sequence of row operations  $op_1$ ,  $op_2$ , . .

**Solution.** Apply the row operations in sequence to the identity matrix of the appropriate size

## Question

Find the matrix implementing the following sequence of elementary row operations on a  $3 \times n$  matrix.

$$R_2 \leftarrow 3R_2$$

$$R_1 \leftarrow R_1 + R_2$$

$$R_2 \leftrightarrow R_3$$

Then multiply it with the all-ones 3×3 matrix.

#### Answer

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 3 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix} \qquad R_2 \leftrightarrow R_3$$

$$R_1 \leftarrow R_1 + R_2$$

$$R_2 \leftrightarrow R_3$$

# Second Key Observation

#### Second Key Observation

Elementary row operations are **invertible** linear transformations

#### Second Key Observation

Elementary row operations are **invertible** linear transformations

This also means the product of elementary matrices is invertible

$$(E_1 E_2 E_3 E_4)^{-1} = E_4^{-1} E_3^{-1} E_2^{-1} E_1^{-1}$$
  
!! the order reverses !!

Describe the inverse transformation for each elementary row operation

Describe the inverse transformation for each elementary row operation

The inverse of scaling by k is scaling by 1/k

Describe the inverse transformation for each elementary row operation

The inverse of scaling by k is scaling by 1/kThe inverse of  $R_i \leftarrow R_i + kR_j$  is  $R_i \leftarrow R_i - kR_j$ 

Describe the inverse transformation for each elementary row operation

The inverse of scaling by k is scaling by 1/k

The inverse of  $R_i \leftarrow R_i + kR_j$  is  $R_i \leftarrow R_i - kR_j$ 

The inverse of swapping is swapping again

#### Recall: Elementary Row Operations

scaling multiply a row by a number

interchange switch two rows

replacement add a scaled equation to another

## Recall: Elementary Row Operations

We only need these two for the forward phase

interchange switch two rows

replacement add a scaled equation to another

#### Recall: Elementary Row Operations

We'll assume we only need this

replacement add a scaled equation to another

#### Reminder: LU Factorization at a High Level

Given a  $m \times n$  matrix A, we are going to factorize A as

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ * & 1 & 0 & 0 \\ * & * & 1 & 0 \\ * & * & * & 1 \end{bmatrix} \begin{bmatrix} \blacksquare & * & * & * & * \\ 0 & \blacksquare & * & * & * \\ 0 & 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$L \qquad U$$

$$A \sim A_1 \sim A_2 \sim \dots \sim A_k$$

Consider a sequence of elementary row operations from  $\boldsymbol{A}$  to an echelon form

Each step can be represent as a product with an elementary matrix

$$A \sim E_1 A \sim E_2 E_1 A \sim \dots \sim E_k E_{k-1} \dots E_2 E_1 A$$

$$A \sim E_1 A \sim E_2 E_1 A \sim \dots \sim E_k E_{k-1} \dots E_2 E_1 A$$

This exactly tells us that if B is the final echelon form we get then

$$B = (E_k E_{k-1} ... E_2 E_1)A = EA$$

where E implements a <u>sequence</u> of row operations. So:

$$A \sim E_1 A \sim E_2 E_1 A \sim \dots \sim E_k E_{k-1} \dots E_2 E_1 A$$

This exactly tells us that if B is the final echelon form we get then 

Invertible

$$B = (E_k E_{k-1} ... E_2 E_1)A = EA$$

where E implements a <u>sequence</u> of row operations. So:

$$A \sim E_1 A \sim E_2 E_1 A \sim \dots \sim E_k E_{k-1} \dots E_2 E_1 A$$

This exactly tells us that if B is the final echelon form we get then 

Invertible

$$B = (E_k E_{k-1} ... E_2 E_1)A = EA$$

where E implements a <u>sequence</u> of row operations. So:

$$A = E^{-1}B = (E_1^{-1}E_2^{-1}...E_{k-1}^{-1}E_k^{-1})B$$

# LU Factorization Algorithm

# LU Factorization Algorithm

1 FUNCTION LU\_Factorization(A):

#### LU Factorization Algorithm

```
1 FUNCTION LU_Factorization(A):
2   L ← identity matrix
```

```
1 FUNCTION LU_Factorization(A):
2    L ← identity matrix
3    U ← A
```

```
1 FUNCTION LU_Factorization(A):
2    L ← identity matrix
3    U ← A
4    convert U to an echelon form by GE forward step # without swaps
```

```
1 FUNCTION LU_Factorization(A):
2    L ← identity matrix
3    U ← A
4    convert U to an echelon form by GE forward step # without swaps
5    FOR each row operation OP in the prev step:
```

```
1 FUNCTION LU_Factorization(A):
2    L ← identity matrix
3    U ← A
4    convert U to an echelon form by GE forward step # without swaps
5    FOR each row operation OP in the prev step:
6    E ← the matrix implementing OP
```

```
FUNCTION LU_Factorization(A):
       L ← identity matrix
       \mathsf{U} \leftarrow A
       convert U to an echelon form by GE forward step # without swaps
       FOR each row operation OP in the prev step:
           E ← the matrix implementing OP
6
           L \leftarrow L @ E^{-1} # note the multiplication on the right
```

```
FUNCTION LU_Factorization(A):
       L ← identity matrix
       \mathsf{U} \leftarrow A
       convert U to an echelon form by GE forward step # without swaps
       FOR each row operation OP in the prev step:
           E ← the matrix implementing OP
6
           L \leftarrow L @ E^{-1} # note the multiplication on the right
       RETURN (L, U)
```

```
FUNCTION LU_Factorization(A):
       L ← identity matrix
       \mathsf{U} \leftarrow A
       convert U to an echelon form by GE forward step # without swaps
       FOR each row operation OP in the prev step:
           E ← the matrix implementing OP
6
           L \leftarrow L @ E^{-1} # note the multiplication on the right
                          we'll see how to do this more efficiently
       RETURN (L, U)
```

# The forward part of Gaussian elimination <u>is</u> matrix factorization

#### The "L" Part

$$E = E_k E_{k-1} \dots E_2 E_1$$

This a product of elementary matrices

So  $L = E^{-1} = E_1^{-1} E_2^{-1} ... E_{k-1}^{-1} E_k^{-1}$  !! the order reverses !!

We won't prove this, but it's worth thinking about: why is this lower triangular?

And can we build this in a more efficient way?

# demo

# How To: LU Factorization by hand

**Question.** Find a LU Factorization for the matrix A (assuming no swaps)

#### Solution.

- $\gg$  Start with L as the identity matrix
- $\gg$  Find U by the forward part of GE
- » For each operation  $R_i \leftarrow R_i + kR_j$ , set  $L_{ij}$  to -k

#### Practice Problem

 1
 2

 3
 7

Determine an LU factorization of the above matrix using this procedure

#### Answer

1237

Determine an LU factorization of the above matrix using this procedure

# Analyzing LU Factorization

We will not use  $O(\cdot)$  notation!

```
We will not use O(\cdot) notation!
```

For numerics, we care about number of **FL**oating-oint **OP**erations (FLOPs):

- >> addition
- >> subtraction
- >> multiplication
- >> division
- >> square root

```
We will not use O(\cdot) notation!
```

For numerics, we care about number of **FL**oating-oint **OP**erations (FLOPs):

- >> addition
- >> subtraction
- >> multiplication
- >> division
- >> square root

```
2n vs. n is very different when n \sim 10^{20}
```

That said, we don't care about exact bounds

That said, we don't care about exact bounds

A function f(n) is asymptotically equivalent to g(n) if

$$\lim_{i \to \infty} \frac{f(i)}{g(i)} = 1$$

That said, we don't care about exact bounds

A function f(n) is asymptotically equivalent to g(n) if

$$\lim_{i \to \infty} \frac{f(i)}{g(i)} = 1$$

For polynomials, they are equivalent to their dominant term

Dominant Terms 
$$\lim_{i \to \infty} \frac{3i^2}{4i^2} \neq 1$$

the dominant term of a polynomial is the monomial with the highest degree

$$\lim_{i \to \infty} \frac{3x^3 + 100000x^2}{3x^3} = 1$$

 $3x^3$  dominates the function even though the coefficient for  $x^2$  is so large

# How To: Solving systems with the LU

**Question.** Solve the equation  $A\mathbf{x} = \mathbf{b}$  given that A = LU is a LU factorization.

**Solution.** First solve  $L\mathbf{x} = \mathbf{b}$  to get a solution  $\mathbf{c}$ , then solve  $U\mathbf{x} = \mathbf{c}$  to get a solution  $\mathbf{d}$ .

Verify: 
$$(Lu)\dot{x}=\dot{b}$$
  $L(u\dot{x})=\dot{b}$   $A\dot{d}=$   
 $L\dot{q}=\dot{b}$   $\Rightarrow$   $\gamma=\dot{c}$   $U\dot{x}=\dot{c}$   $x=\dot{d}$   $L\dot{c}=\dot{b}$ 

# How To: Solving systems with the LU

**Question.** Solve the equation  $A\mathbf{x} = \mathbf{b}$  given that A = LU is a LU factorization.

**Solution.** First solve  $L\mathbf{x} = \mathbf{b}$  to get a solution  $\mathbf{c}$ , then solve  $U\mathbf{x} = \mathbf{c}$  to get a solution  $\mathbf{d}$ .

Why is this better than just solving Ax = b?

The following FLOP estimates are based on  $n \times n$  matrices

The following FLOP estimates are based on  $n \times n$  matrices

Gaussian Elimination: 
$$\sim \frac{2n^3}{3}$$
 FLOPS

The following FLOP estimates are based on  $n \times n$  matrices

Gaussian Elimination: 
$$\sim \frac{2n^3}{3}$$
 FLOPS

GE Forward:  $\sim \frac{2n^3}{3}$  FLOPS

The following FLOP estimates are based on  $n \times n$  matrices

Gaussian Elimination: 
$$\sim \frac{2n^3}{3}$$
 FLOPS

GE Forward:  $\sim \frac{2n^3}{3}$  FLOPS

GE Backward:  $\sim 2n^2$  FLOPS

The following FLOP estimates are based on  $n \times n$  matrices

Gaussian Elimination:  $\sim \frac{2n^3}{3}$  FLOPS

GE Forward:  $\sim \frac{2n^3}{3}$  FLOPS

GE Backward:  $\sim 2n^2$  FLOPS

Matrix Inversion:  $\sim 2n^3$  FLOPS

The following FLOP estimates are based on  $n \times n$  matrices

Gaussian Elimination:  $\sim \frac{2n^3}{3}$  FLOPS

GE Forward:  $\sim \frac{2n^3}{3}$  FLOPS

GE Backward:  $\sim 2n^2$  FLOPS

Matrix Inversion:  $\sim 2n^3$  FLOPS

Matrix-Vector Multiplication:  $\sim 2n^2$  FLOPS

The following FLOP estimates are based on  $n \times n$  matrices

Gaussian Elimination:  $\sim \frac{2n^3}{3}$  FLOPS

GE Forward:  $\sim \frac{2n^3}{3}$  FLOPS

GE Backward:  $\sim 2n^2$  FLOPS

Matrix Inversion:  $\sim 2n^3$  FLOPS

Matrix-Vector Multiplication:  $\sim 2n^2$  FLOPS

Solving by matrix inversion:  $\sim 2n^3$  FLOPS

The following FLOP estimates are based on  $n \times n$  matrices

Gaussian Elimination:  $\sim \frac{2n^3}{3}$  FLOPS

GE Forward:  $\sim \frac{2n^3}{3}$  FLOPS

GE Backward:  $\sim 2n^2$  FLOPS

Matrix Inversion:  $\sim 2n^3$  FLOPS

Matrix-Vector Multiplication:  $\sim 2n^2$  FLOPS

Solving by matrix inversion:  $\sim 2n^3$  FLOPS

Solving by Gaussian elimination:  $\sim \frac{2n^3}{3}$  FLOPS

# FLOPS for solving LU systems

LU Factorization: 
$$\sim \frac{2n^3}{3}$$
 FLOPS Solving  $L\mathbf{x} = \mathbf{b}$ :  $\sim 2n^2$  FLOPS (by "forward" elimination)

Solving  $U\mathbf{x} = \mathbf{c}$ :  $\sim 2n^2$  FLOPS (already in echelon form)


Solving by LU Factorization:  $\sim \frac{2n^3}{2}$  FLOPS

If you solve several matrix equations for the same matrix, LU factorization is <u>faster</u> than matrix inversion on the first equation, and the same (asymptotically) in later equations (and it works for rectangular matrices).

If A doesn't have to many entries (A is **sparse**), then its likely that L and U won't either.

If A doesn't have to many entries (A is **sparse**), then its likely that L and U won't either.

But  $A^{-1}$  may have *many* entries  $(A^{-1}$  is dense)



If A doesn't have to many entries (A is sparse), then its likely that L and U won't either.


But  $A^{-1}$  may have *many* entries  $(A^{-1}$  is dense)

Sparse matrices are faster to compute with and better with respect to storage.

## Algebraic Graph Theory


#### Graphs

**Definition (Informal).** A **graph** is a collection of nodes with edges between them.




#### Directed vs. Undirected Graphs

A graph is **directed** if its edges have a direction.




#### Weighted vs Unweighted graphs


A graph is weighted if its edges have associated values.



#### Weighted vs Unweighted graphs


A graph is weighted if its edges have associated values.





#### Simple Graphs

A graph is **simple** if it is undirected, has no self loops, and no multi-edges.



#### Four Kinds of Graphs

directed

undirected

weighted

nodes are traffic lights
edges are streets
weights are number of lanes

nodes are musicians edges are collaborations weights are number of collaborations

unweighted

nodes are instagram users edges are follows

nodes are bodies of land edges are pedestrian bridges

#### Four Kinds of Graphs

directed

undirected

weighted

nodes are traffic lights
edges are streets
weights are number of lanes

nodes are musicians edges are collaborations weights are number of collaborations

unweighted

nodes are instagram users edges are follows

nodes are bodies of land edges are pedestrian bridges

Today

#### Four Kinds of Graphs

directed undirected

weighted

nodes are traffic lights
edges are streets
weights are number of lanes

Markov Chains

nodes are musicians edges are collaborations weights are number of collaborations

unweighted

nodes are instagram users edges are follows

nodes are bodies of land edges are pedestrian bridges

Today

#### Fundamental Question

#### Fundamental Question

How do we represent a graph formally in a computer?

#### Fundamental Question

How do we represent a graph formally in a computer?


There are a couple ways, but one way is to use <u>matrices</u>.


#### Adjacency Matrices

Let G be an simple graph with its nodes labeled by numbers 1 through  $n_{\bullet}$ 

We can create the adjacency matrix A for G as follows.

$$A_{ij} = \begin{cases} 1 & \text{there is an edge between i and j} \\ 0 & \text{otherwise} \end{cases}$$





#### Symmetric Matrices

**Definition.** A  $n \times n$  matrix is symmetric if

$$A^T = A$$

Example.

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}$$


# Once we have an adjacency matrix, we can do linear algebra on graphs.

Given an adjacency matrix A, can we interpret anything meaningful from  $A^2$ ?

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}$$


$$(A^2)_{53} = 1(0) + 1(1) + 0(0) + 1(1) + 0(0) + 0(0) = 2$$

$$(A^2)_{ij} = A_{i1}A_{1j} + A_{i2}A_{2j} + \dots + A_{in}A_{nj}$$




$$(A^2)_{ij} = A_{i1}A_{1j} + A_{i2}A_{2j} + \dots + A_{in}A_{nj}$$

$$A_{ik}A_{kj} = \begin{cases} 1 & \text{there are edges i to k and k to j} \\ 0 & \text{otherwise} \end{cases}$$




$$(A^2)_{ij} = A_{i1}A_{1j} + A_{i2}A_{2j} + \dots + A_{in}A_{nj}$$

$$A_{ik}A_{kj} = \begin{cases} 1 & \text{there are edges i to k and k to j} \\ 0 & \text{otherwise} \end{cases}$$




$$(A^2)_{ij} = A_{i1}A_{1j} + A_{i2}A_{2j} + \dots + A_{in}A_{nj}$$

 $(A^2)_{ij} = \begin{bmatrix} \text{number of 2-step paths} \\ \text{from i to j} \end{bmatrix}$ 



A **triangle** in an undirected graph is a set of three distinct nodes with edges between every pair of nodes.

Triangles in a social network represent mutual friends and tight cohesion (among other things)



#### Application: Triangle Counting (Naive)

```
FUNCTION tri_count_naive(A):
  count = 0
  for i from 1 to n:
    for j from i + 1 to n:
      for k from j + 1 to n:
        if A_{ij}=1 and A_{jk}=1 and A_{ki}=1: # an edge between each pair
           count += 1:
  RETURN count
```

**Theorem.** For an adjacency matrix A, the number of triangle containing the edge (i,j) is

$$(A^2)_{ij} * A_{ij}$$

Verify:

```
FUNCTION tri_count(A):

compute A^2

count \leftarrow sum of (A^2)_{ij} * A_{ij} for all distinct i and j

RETURN count / 6 # why divided by 6?
```

```
FUNCTION tri_count(A):
 # in NumPy '*' is entry—wise multiplication
        also called the HADAMARD PRODUCT
 #
  count \leftarrow sum of the entries of A^2 * A
  RETURN count / 6
```

```
FUNCTION tri_count(A):
 # in NumPy '*' is entry-wise multiplication
        also called the HADAMARD PRODUCT
 #
 # and 'np.sum' sums the entry of a matrix
 RETURN np.sum((A @ A) * A) / 6
```


### demo

#### Another Application: Reachability

**Question:** If  $A^2$  gives us information about length 2 paths, then what about  $A^k$ ?


 $A^k$  gives us information about k-length paths.

#### Example



#### Example


$$\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix} = \text{adjacency matrix for } G$$



#### Example

$$\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix} = adjacency matrix for G$$


$$\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}^2 = \begin{bmatrix} 2 & 0 & 0 & 2 \\ 0 & 2 & 2 & 0 \\ 0 & 2 & 2 & 0 \\ 2 & 0 & 0 & 2 \end{bmatrix}$$




$$egin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix} = adjacency matrix for  $G$$$


$$\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}^{2} = \begin{bmatrix} 2 & 0 & 0 & 2 \\ 0 & 2 & 2 & 0 \\ 0 & 2 & 2 & 0 \\ 2 & 0 & 0 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}^{3} = \begin{bmatrix} 2 & 0 & 0 & 2 \\ 0 & 2 & 2 & 0 \\ 0 & 2 & 2 & 0 \\ 2 & 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 4 & 4 & 0 \\ 4 & 0 & 0 & 4 \\ 4 & 0 & 0 & 4 \\ 0 & 4 & 4 & 0 \end{bmatrix}$$






```
\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} = adjacency matrix for G
```



$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} = adjacency matrix for G$$


$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}^2 = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \\ 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$



$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} = adjacency matrix for G$$

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}^2 = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \\ 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$


$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}^{3} = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \\ 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 2 & 0 & 1 \\ 2 & 0 & 3 & 0 \\ 0 & 3 & 0 & 2 \\ 1 & 0 & 2 & 0 \end{bmatrix}$$




### Another Application: Reachability

Theorem: Let G be a simple graph.

- $(A_G^k)_{ij}$  is the number of paths of length exactly k from  $v_i$  to  $v_i$ .
- $\left((A_G+I)^k\right)_{ij}$  is nonzero if and only if there is a path of length at at most k from  $v_i$  to  $v_j$ .




$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix} = (adjacency matrix for  $G) + I$$$



$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix} = (adjacency matrix for  $G) + I$$$


$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix}^2 = \begin{bmatrix} 3 & 2 & 2 & 2 \\ 2 & 3 & 2 & 2 \\ 2 & 2 & 3 & 2 \\ 2 & 2 & 2 & 3 \end{bmatrix}$$



$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix} = (adjacency matrix for  $G) + I$$$

$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix}^2 = \begin{bmatrix} 3 & 2 & 2 & 2 \\ 2 & 3 & 2 & 2 \\ 2 & 2 & 3 & 2 \\ 2 & 2 & 2 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}^{3} = \begin{bmatrix} 3 & 2 & 2 & 2 \\ 2 & 3 & 2 & 2 \\ 2 & 2 & 3 & 2 \\ 2 & 2 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 6 & 6 & 6 & 7 \\ 6 & 6 & 7 & 6 \\ 6 & 7 & 6 & 6 \\ 7 & 6 & 6 & 6 \end{bmatrix}$$




### How To: Reachability

**Question:** Given a simple graph G determine how many nodes  $v_i$  can reach in at least k steps.

**Answer:** Find  $(A_G + I)^k$  and count the number of nonzero elements in column i.

## Question

Determine the  $(A_G+I)^2$  and  $(A_G+I)^3$  and interpret the results.



#### Summary

Matrix inverses allow us to easily solve many matrixes equations over the same  $\boldsymbol{A}$ 

LU Factorizations allows us to do the same, but more generally more efficiently

Adjacency matrices are linear algebraic representations of graphs