

Analytic Geometry in \mathbb{R}^n

Geometric Algorithms

Lecture 21

Practice Problem

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

$$\left\{ \begin{bmatrix} 0 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

can't do in general,

Let A be a 4×4 matrix with eigenvalues 3 and -2 where $\dim(\text{Nul}(A + 2I)) = 3$.

but you can with
eigenvectors from distinct
eigenspaces

True or False: A must be diagonalizable.

3 lin ind. eigenvectors from -2 eigenspace

≥ 1 eigenvectors from 3 eigenspace

\rightarrow 4 lin ind. eigenvectors

Answer: True

The set of eigenvectors we get from the diagonalization procedure is of size 4, which means there is an eigenbasis of \mathbb{R}^4 for A .

Objectives

1. Recall what we learned in algebra class
2. Connect the familiar notions of lengths, distances, and angles to inner products
3. Begin discussing the fundamental concept of orthogonality

Keywords


inner product

norm

orthogonal


Motivation

What is Analytic Geometry?



What is Analytic Geometry?


Analytic geometry is the study of space using a coordinate system.



What is Analytic Geometry?

Analytic geometry is the study of space using a coordinate system.

We're interested in equations about lines, curves, shapes, angles, etc.




What is Analytic Geometry?

Analytic geometry is the study of space using a coordinate system.

We're interested in equations about lines, curves, shapes, angles, etc.

The fundamental concepts are:




What is Analytic Geometry?

Analytic geometry is the study of space using a coordinate system.


We're interested in equations about lines, curves, shapes, angles, etc.

The fundamental concepts are:

- » distance
- » position
- » area
- » angle




A Potentially Familiar Example




What is the value of θ ?

Angles in \mathbb{R}^2



What is the value of θ ?

Angles in \mathbb{R}^3




What is the value of θ ?

The First Key Idea

Angles make sense in *any* dimension.



Any pair of vectors in \mathbb{R}^n span a (2D) plane.

(We could formalize this via change of bases)



The Picture

We can do "normal" analytic geometry here



change of basis from $\text{span}\{v, w\}$ to \mathbb{R}^2
(caveat: doesn't always preserve geometry)

A Fundamental Question

Doing this change of basis every time we want to do geometry is a lot of work...

Can we do it directly using ideas we've been learning?

Recall: Inner Products

Recall: Inner Products

$$[u_1 \quad u_2 \quad u_3 \quad u_4] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = u_1 v_1 + u_2 v_2 + u_3 v_3 + u_4 v_4$$

Recall: Inner Products

$$[u_1 \quad u_2 \quad u_3 \quad u_4] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = u_1 v_1 + u_2 v_2 + u_3 v_3 + u_4 v_4$$

Definition. The **inner product** of two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n is

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v}$$

Recall: Inner Products

$$[u_1 \ u_2 \ u_3 \ u_4] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = u_1 v_1 + u_2 v_2 + u_3 v_3 + u_4 v_4$$

Definition. The **inner product** of two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n is **a.k.a. dot product**

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v}$$

standard inner product
on \mathbb{R}^n

The Second Key Idea

The Second Key Idea

All of the basic concepts of analytic geometry
can be defined *in terms of inner products*.

The Second Key Idea

All of the basic concepts of analytic geometry can be defined *in terms of inner products*.

Definition (Advanced). An **inner product space** is a vector space with an inner product function.

The Second Key Idea

All of the basic concepts of analytic geometry can be defined *in terms of inner products*.

Definition (Advanced). An **inner product space** is a vector space with an inner product function.

Inner product spaces (like \mathbb{R}^n) are places where you can do analytic geometry.

The Fundamental Question

How do we do analytic geometry,
given we have an inner product?

Inner Products

Recall: Inner Products (Again)

$$[u_1 \ u_2 \ u_3 \ u_4] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = u_1 v_1 + u_2 v_2 + u_3 v_3 + u_4 v_4$$

Definition. The **inner product** of two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n is **a.k.a. dot product**

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v}$$

Example

$$\mathbf{u} = \begin{bmatrix} -3 \\ 2 \\ 1 \\ 4 \end{bmatrix}$$

$$\mathbf{v} = \begin{bmatrix} 2 \\ 0.5 \\ -1 \\ 3 \end{bmatrix}$$

$$\langle \vec{u}, \vec{v} \rangle = \vec{u}^T \vec{v} = -6 + 1 - 1 + 12 = 6$$

Algebraic Properties of Inner Products

- $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$ (symmetry)

$$(A+B)\vec{v} = A\vec{v} + B\vec{v}$$

- $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = (\mathbf{u} \cdot \mathbf{w}) + (\mathbf{v} \cdot \mathbf{w})$
- $(\alpha\mathbf{u}) \cdot \mathbf{v} = \alpha(\mathbf{u} \cdot \mathbf{v})$

} linearity in the first argument


- $\mathbf{u} \cdot \mathbf{u} \geq 0$ (nonnegativity)

- $\mathbf{u} \cdot \mathbf{u} = 0$ if and only if $\mathbf{u} = 0$

} positive-definite

Verifying Additivity

$$\begin{aligned}\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle &= (\vec{u} + \vec{v})^T \vec{w} \\ &= (\vec{u}^T + \vec{v}^T) \vec{w} \\ &= \vec{u}^T \vec{w} + \vec{v}^T \vec{w}\end{aligned}$$


$$= \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$$

Homogeneity in the Right Argument

$$\langle \mathbf{v}, c\mathbf{u} \rangle = c \langle \mathbf{v}, \mathbf{u} \rangle$$

Verify:

$$\vec{v}^T \overbrace{c\vec{u}}^{//} = c \overbrace{\vec{v}^T \vec{u}}^{//}$$

An Aside: What is this linear transformation?

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \mapsto \begin{bmatrix} 3 \\ 5 \\ 7 \end{bmatrix} \cdot \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \quad \begin{matrix} \mathbb{R}^3 \rightarrow \mathbb{R} \\ 1 \times 3 \end{matrix}$$

Let's find the matrix for this transformation:

$$1 \cdot \{ \overbrace{[3 \ 5 \ 7]}^3 \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}$$

Algebraic Properties of Inner Products

- $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$ (symmetry)
 - $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = (\mathbf{u} \cdot \mathbf{w}) + (\mathbf{v} \cdot \mathbf{w})$
 - $(\alpha \mathbf{u}) \cdot \mathbf{v} = \alpha(\mathbf{u} \cdot \mathbf{v})$
- } linearity in the first argument
- $\mathbf{u} \cdot \mathbf{u} \geq 0$ (nonnegativity)
 - $\mathbf{u} \cdot \mathbf{u} = 0$ if and only if $\mathbf{u} = \mathbf{0}$

Nonnegativity

$$\langle \mathbf{v}, \mathbf{v} \rangle = \sum_{i=1}^n v_i^2$$

Nonnegativity

$$\langle \mathbf{v}, \mathbf{v} \rangle = \sum_{i=1}^n v_i^2$$

Squared values are always nonnegative.

Nonnegativity

$$\langle \mathbf{v}, \mathbf{v} \rangle = \sum_{i=1}^n v_i^2 \stackrel{\text{if}}{=} 0 \Rightarrow v_i^2 = 0 \text{ for all } i$$

Squared values are always nonnegative. $\Rightarrow v_i = 0$ for all i

Therefore $\langle \mathbf{v}, \mathbf{v} \rangle$ is always nonnegative.

Nonnegativity

$$\langle \mathbf{v}, \mathbf{v} \rangle = \sum_{i=1}^n v_i^2$$

Squared values are always nonnegative.

Therefore $\langle \mathbf{v}, \mathbf{v} \rangle$ is always nonnegative.

Question. What happens when we scale a vector to make it longer?

Nonnegativity and Scaling

$$\langle c\mathbf{v}, c\mathbf{v} \rangle = c^2 \langle \mathbf{v}, \mathbf{v} \rangle = c^2 \sum_{i=1}^n v_i^2$$

Nonnegativity and Scaling

$$\langle c\mathbf{v}, c\mathbf{v} \rangle = c^2 \langle \mathbf{v}, \mathbf{v} \rangle = c^2 \sum_{i=1}^n v_i^2$$

If $c > \cancel{0}$ then $\langle c\mathbf{v}, c\mathbf{v} \rangle > \langle \mathbf{v}, \mathbf{v} \rangle$.

Nonnegativity and Scaling

$$\langle c\mathbf{v}, c\mathbf{v} \rangle = c^2 \langle \mathbf{v}, \mathbf{v} \rangle = c^2 \sum_{i=1}^n v_i^2$$

If $c > 0$ then $\langle c\mathbf{v}, c\mathbf{v} \rangle > \langle \mathbf{v}, \mathbf{v} \rangle$.

Increasing the length of a vector increases its inner product with itself.

Nonnegativity and Scaling

$$\langle c\mathbf{v}, c\mathbf{v} \rangle = c^2 \langle \mathbf{v}, \mathbf{v} \rangle = c^2 \sum_{i=1}^n v_i^2$$

If $c > 0$ then $\langle c\mathbf{v}, c\mathbf{v} \rangle > \langle \mathbf{v}, \mathbf{v} \rangle$.

Increasing the length of a vector increases its inner product with itself.

This means $\langle \mathbf{v}, \mathbf{v} \rangle$ is capturing some notion of magnitude.

The Fundamental Question

How does this all connect back to
distances and angles?

Question

$$(a+b)(a-b) \\ = a^2 - b^2$$

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$$

$$(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = (\mathbf{u} \cdot \mathbf{w}) + (\mathbf{v} \cdot \mathbf{w})$$

Simplify the expression $\langle \mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle$ using the properties of inner products.

$$= (\vec{u} + \vec{v})^T (\vec{u} - \vec{v}) = (\vec{u} + \vec{v})^T \vec{u} - (\vec{u} + \vec{v})^T \vec{v}$$


$$= \vec{u}^T \vec{u} + \vec{v}^T \vec{u} - \vec{u}^T \vec{v} - \vec{v}^T \vec{v}$$

↖ ↗
same

Answer: $\langle \mathbf{u}, \mathbf{u} \rangle - \langle \mathbf{v}, \mathbf{v} \rangle$

Norms (Lengths/Distances)

Another Potentially Familiar Question




$$\begin{aligned} c^2 &= 3^2 + 4^2 \\ &= 9 + 16 \\ &= 25 \end{aligned}$$

$$\Rightarrow c = 5$$

How long is the side c ?

Pythagorean Theorem



Theorem (Pythagoras). *For a right triangle, the square of the length of the hypotenuse is the sum of the squares of the lengths of the remaining two sides.*

This still works in \mathbb{R}^3


Theorem (Pythagoras). $C = \sqrt{v_1^2 + v_2^2 + v_3^2}$

Verify:

$$C = \sqrt{G^2 + v_3^2}$$

$$G = \sqrt{v_1^2 + v_2^2}$$

$$C = \sqrt{v_1^2 + v_2^2 + v_3^2}$$



Norm

Definition. The (ℓ^2) norm of a vector \mathbf{v} in \mathbb{R}^n is

$$\|\mathbf{v}\| = \left\| \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \right\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2} = \sqrt{\sum_{i=1}^n v_i^2}$$

The norm of a vector is the square root of the sum of the squares of its entries.

Norms and Inner Products

Definition. The ℓ^2 norm of a vector \mathbf{v} in \mathbb{R}^n is

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}}$$

The norm of a vector is the square root of the inner product with itself.

Norms and Inner Products

Definition. The ℓ^2 norm of a vector \mathbf{v} in \mathbb{R}^n is

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}}$$


The norm of a vector is the square root of the inner product with itself.

It's important that $\mathbf{v}^T \mathbf{v}$ is nonnegative.

Norms and Distance

Norms give us a notion of length.

In \mathbb{R}^2 and \mathbb{R}^3 this is our existing notion of length.




ℓ^2 Normalization

Definition. A **unit vector** is a vector \mathbf{v} such that $\|\mathbf{v}\| = 1$.

We often *normalize* vectors if we only care about their direction:

$$\mathbf{v} \mapsto \frac{\mathbf{v}}{\|\mathbf{v}\|}$$

$$\frac{\vec{v}}{\|\vec{v}\|} \cdot \frac{\vec{v}}{\|\vec{v}\|} = \frac{\vec{v}^T \vec{v}}{\|\vec{v}\|^2} = \frac{\|\vec{v}\|^2}{\|\vec{v}\|^2} = 1$$



How To: Normalizing Vectors

Question. Find the unit vector which points in the same direction as \mathbf{u} .

Solution. Compute $\|\mathbf{u}\|$. The unit vector is then

$$\frac{\mathbf{u}}{\|\mathbf{u}\|}$$

Example

Find the unit vector in the same direction as $\vec{v} = \begin{bmatrix} 1 \\ -2 \\ 2 \\ 0 \end{bmatrix}$

$$\begin{aligned} \|\vec{v}\| &= \sqrt{1^2 + (-2)^2 + (2)^2} \\ &= \sqrt{1 + 4 + 4} = \sqrt{9} = 3 \end{aligned}$$

(opposite)

$$\frac{\vec{v}}{\|\vec{v}\|} =$$

$$\begin{bmatrix} 1/3 \\ -2/3 \\ 2/3 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -1/3 \\ 2/3 \\ -2/3 \\ 0 \end{bmatrix}$$


The Unit Sphere





Definition. The unit n -sphere is the collection of all unit vectors in \mathbb{R}^n .

Vector norms allow us to talk about spheres in higher dimensions.

A sphere is a collection of points equidistant from a center point.





An Aside: Other Notions of Distance



An Aside: Other Notions of Distance



Why are we talking about norms and inner products so generally?



An Aside: Other Notions of Distance

Why are we talking about norms and inner products so generally?

Because there are other inner products and norms.





An Aside: Other Notions of Distance

Why are we talking about norms and inner products so generally?

Because there are other inner products and norms.

e.g., Manhattan distance




Another Aside: Surface Area and Volume

(curse of dimensionality)

With a bit of calculus, we can calculate the surface area and volume of the unit n -sphere.

And the result is bizarre...

the infinite dimensional unit sphere has no volume or surface area...




moving on...

Distance

If we know how to calculate
lengths of vectors, we know how
to calculate distances.

Recall: Vector Addition


tip-to-tail rule:



Recall: Vector Addition

tip-to-tail rule:


$\mathbf{u} + \mathbf{v}$ result of putting
the tail of \mathbf{v} to the tip
of \mathbf{u} (or vice versa)



Recall: Vector Addition

tip-to-tail rule:


$\mathbf{u} + \mathbf{v}$ result of putting
the tail of \mathbf{v} to the tip
of \mathbf{u} (or vice versa)



Recall: Vector Addition

tip-to-tail rule:

$\mathbf{u} + \mathbf{v}$ result of putting
the tail of \mathbf{v} to the tip
of \mathbf{u} (or vice versa)




Recall: Vector Addition



The distance between u and $u+v$ is the length of v

tip-to-tail rule:

$u+v$ result of putting the tail of v to the tip of u (or vice versa)



Distance (Pictorially)



$$\vec{v} + (\vec{u} - \vec{v}) = \vec{u}$$

$$\text{dist}(\vec{u}, \vec{v}) = \|\vec{u} - \vec{v}\|$$

Distance (Algebraically)


Definition. The distance between two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n is given by

$$\text{dist}(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\| = \|\vec{u} - \vec{v}\|$$

e.g., $\mathbf{u} = \begin{bmatrix} 7 \\ 1 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$

$$\vec{u} - \vec{v} = \begin{bmatrix} 4 \\ -1 \end{bmatrix} \Rightarrow \|\vec{u} - \vec{v}\| = \sqrt{4^2 + (-1)^2} = \sqrt{17}$$

Question



$$d = \text{dist}\left(\frac{\vec{v}}{\|\vec{v}\|}, \frac{\vec{u}}{\|\vec{u}\|}\right)$$

$$= \left\| \frac{\vec{v}}{\|\vec{v}\|} - \frac{\vec{u}}{\|\vec{u}\|} \right\|$$

$$d' = \text{dist}\left(\frac{2\vec{v}}{\|2\vec{v}\|}, \frac{2\vec{u}}{\|2\vec{u}\|}\right)$$

$$= \left\| \frac{2\vec{v}}{\|2\vec{v}\|} - \frac{2\vec{u}}{\|2\vec{u}\|} \right\|$$

Find an expression for the distance d .

$$= 2 \left\| \frac{\vec{v}}{\|\vec{v}\|} - \frac{\vec{u}}{\|\vec{u}\|} \right\|$$

Challenge. Find an expression for d' .


$$= 2d$$

Answer

Angles

Again, Angles still make sense


Any pair of vectors in \mathbb{R}^n
span a (2D) plane.



Fundamental Question


How do we determine the angle
between any two vectors?

Recall: A Potentially Familiar Example



What is the value of θ ?


Law of Cosines



Theorem.

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$

Law of Cosines




Theorem.

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$

Generalized the Pythagorean Theorem

Law of Cosines




Theorem.

θ exactly when $\theta = 90^\circ$

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$

Generalized the Pythagorean Theorem

The Picture



In more "vector"-y terms:

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2\|\mathbf{u}\|\|\mathbf{v}\|\cos\theta$$

lengths
↓ ↓

Isolating θ

$$\cos \theta = \frac{c^2 - a^2 - b^2}{2ab}$$

$$\theta = \cos^{-1} \left(\frac{c^2 - a^2 - b^2}{2ab} \right)$$

We might remember these equations...

Isolating θ

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2\|\mathbf{u}\|\|\mathbf{v}\|\cos\theta$$

Let's isolate θ in this equation: $\langle \vec{u} - \vec{v}, \vec{u} - \vec{v} \rangle$

$$\begin{aligned} 2\|\vec{u}\|\|\vec{v}\|\cos\theta &= \|\vec{u}\|^2 + \|\vec{v}\|^2 - \|\vec{u} - \vec{v}\|^2 \\ &= \cancel{\|\vec{u}\|^2} + \cancel{\|\vec{v}\|^2} - (\cancel{\|\vec{u}\|^2} + \cancel{\|\vec{v}\|^2} - 2\langle \vec{u}, \vec{v} \rangle) \\ &= 2\langle \vec{u}, \vec{v} \rangle \end{aligned}$$

$$\cos\theta = \frac{\langle \vec{u}, \vec{v} \rangle}{\|\vec{u}\|\|\vec{v}\|}$$

Cosines and Unit Vectors

Theorem. For vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n with an angle θ between them,

$$\cos \theta = \left\langle \frac{\mathbf{u}}{\|\mathbf{u}\|}, \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\rangle = \frac{\langle \vec{u}, \vec{v} \rangle}{\|\vec{u}\| \|\vec{v}\|}$$

The cosine of the angle between two vectors is the inner product of their ℓ^2 normalizations.

How To: Angles

Question. Find the angle between the two vectors \mathbf{u} and \mathbf{v} .

Solution. Compute $\cos^{-1} \left(\frac{\mathbf{u}}{\|\mathbf{u}\|} \cdot \frac{\mathbf{v}}{\|\mathbf{v}\|} \right)$ (with a calculator).

Example

Find the angle between the vectors

$$\mathbf{u} = \begin{bmatrix} 1 \\ 3 \\ -7 \\ -2 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} 8 \\ -2 \\ 4 \\ 6 \end{bmatrix}$$

Example: Step 1

Compute $\|\mathbf{u}\|$ and $\|\mathbf{v}\|$.

$$\|\mathbf{u}\| = \sqrt{1^2 + 3^2 + (-7)^2 + (-2)^2} = 7.93$$

$$\|\mathbf{v}\| = \sqrt{8^2 + (-2)^2 + 4^2 + 6^2} = 10.95$$

Example: Step 2

Normalize the vectors.

$$\frac{\mathbf{u}}{\|\mathbf{u}\|} = \begin{bmatrix} 0.13 \\ 0.38 \\ -0.88 \\ -0.25 \end{bmatrix}$$

$$\frac{\mathbf{v}}{\|\mathbf{v}\|} = \begin{bmatrix} 0.73 \\ -0.18 \\ 0.36 \\ 0.54 \end{bmatrix}$$

Example: Step 3

Find their inner product.

$$\left\langle \frac{\mathbf{u}}{\|\mathbf{u}\|}, \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\rangle = (0.13 \cdot 0.73) + (0.38 \cdot -0.18) + (-0.88 \cdot 0.36) + (-0.25 \cdot 0.54) \\ = -0.44$$

Example: Step 4

Compute the angle.

$$\theta = \cos^{-1}(-0.44) = 116^\circ$$

Orthogonality (Perpendicularity)

A Simpler Fundamental Question

How do we determine if angle
between any two vectors is 90° ?

Orthogonality

Orthogonality

Definition (Informal). Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are orthogonal if then angle between them is 90° .

Orthogonality

Definition (Informal). Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if then angle between them is 90° .

This isn't actually that informal, it's perfectly reasonable for the purposes of this course.

Orthogonality

Definition (Informal). Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if then angle between them is 90° .

This isn't actually that informal, it's perfectly reasonable for the purposes of this course.

Orthogonal and perpendicular are the same thing.

Orthogonality

Definition (Informal). Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if then angle between them is 90° .

This isn't actually that informal, it's perfectly reasonable for the purposes of this course.

Orthogonal and perpendicular are the same thing.

But it doesn't connect back to inner products.

Orthogonality

Definition (Informal). Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if then angle between them is 90° .

This isn't actually that informal, it's perfectly reasonable for the purposes of this course.

Orthogonal and perpendicular are the same thing.

But it doesn't connect back to inner products.

(and it's difficult to compute with)

Recall: Cosines and Unit Vectors

Theorem. For vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n with an angle θ between them, if $\theta = 90^\circ$ or $\pi/2$

$$\cos \theta = \left\langle \frac{\mathbf{u}}{\|\mathbf{u}\|}, \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\rangle = 0 = \frac{\langle \vec{u}, \vec{v} \rangle}{\|\mathbf{u}\| \|\mathbf{v}\|}$$

The cosine of the angle between two vectors is the inner product of their ℓ^2 normalizations.

Orthogonality


Definition (Actual). Vectors \mathbf{u} and \mathbf{v} are **orthogonal** if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

This definition gives an easy computational way to determine orthogonality.


Example.

$$\begin{bmatrix} 1 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 2 \end{bmatrix} = 2 - 2 = 0$$


Derivation by Picture



Derivation by Picture



Derivation by Picture



Derivation by Algebra

u and **v** are orthogonal exactly when

$$\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 = \|\mathbf{u} - \mathbf{v}\|^2 = \|\vec{u}\|^2 + \|\vec{v}\|^2 - 2\langle \vec{u}, \vec{v} \rangle$$

Let's simplify this a bit:

$$\Downarrow$$
$$\langle \vec{u}, \vec{v} \rangle = 0$$


How To: Orthogonality

Question. Determine if \mathbf{u} and \mathbf{v} are perpendicular.

Solution. Determine if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$. If yes, then they are perpendicular. If no, then they are not.

Application: Cosine Similarity

High Level



Data points are very big vectors.

Similar vectors "point in nearly the same direction."

Example: Netflix Users


$$\text{user}_1 = \begin{bmatrix} 2 \\ 10 \\ 1 \\ 3 \end{bmatrix} \quad \text{user}_2 = \begin{bmatrix} 2 \\ 5 \\ 0 \\ 4 \end{bmatrix} \quad \text{user}_3 = \begin{bmatrix} 10 \\ 0 \\ 5 \\ 6 \end{bmatrix} \quad \begin{array}{l} \text{comedy} \\ \text{drama} \\ \text{horror} \\ \text{romance} \end{array}$$

A Netflix user might be represented as a vectors whose i th entry is the number of movies they've watched in a particular genre.

Who are more likely to share similar interests in movies?

Cosine Similarity

When is $\cos \theta = -1$



Definition. The **cosine similarity** of two vectors is the cosine of the angle between them.

If its close to 0, then two Netflix users watch very different movies.

If its close to 1, then two Netflix users watch very similar movies.


Example: Netflix Users

comedy

drama

horror

romance



Other Examples

- *Document similarity*
 - Documents \mapsto word count vectors
 - Similar documents should use similar words
- *Word2Vec*
 - Words \mapsto vector *somehow*
 - This underlies modern natural language processing (NLP)

Summary

We can talk about distances and angles in \mathbb{R}^n .

Every basic geometric concept connects to inner products.

Once we can talk about distances and angles we can talk about similarity.